



















- Raccuia SA, Piscioneri I, Sharma N, Melilli MG, 2011. Genetic variability in *Cynara cardunculus* L. domestic and wild types for grain oil production and fatty acids composition. *Biomass Bioener.* 35:3167-73.
- Rad A, Abbasian A, Aminpanah H, 2014. Seed and oil yield of rapeseed (*Brassica napus* L.) cultivars under irrigated and non-irrigated condition. *J. Anim. Plant Sci* 24:204-10.
- Sharma N, Hotte N, Rahman MH, Mohammadi M, Deyholos MK, Kav NN, 2008. Towards identifying Brassica proteins involved in mediating resistance to *Leptosphaeria maculans*: a proteomics-based approach. *Proteomics* 8:3516-35.
- UNESCO-FAO, 1963. Bioclimatic map of the Mediterranean zone; NS162/III, 22A. UNESCO, Paris - FAO, Rome.
- Velasco L, Fernández-Martínez JM, De Haro A, 1997. Determination of the fatty acid composition of the oil in intact-seed mustard by near-infrared reflectance spectroscopy. *J. Am. Oil Chem. Soc.* 74:1595-602.
- Velasco L, Nabloussi A, De Haro A, Fernández-Martínez JM, 2003. Development of high-oleic, low-linolenic acid Ethiopian-mustard (*Brassica carinata*) germplasm. *Theor. Appl. Genet.* 107:823-30.
- Wang L, Weller CL, 2006. Recent advances in extraction of nutraceuticals from plants. *Trends Food Sci. Tech.* 17:300-12.
- Zapata N, Vergas M, Reyes JF, Belmar G, 2012. Quality of biodiesel and press cake obtained from *Euphorbia lathyris*, *Brassica napus* and *Ricinus communis*. *Ind. Crop. Prod.* 38:1-5.

Non commercial use only